

Using the local density approximation and the LYP, BLYP, and B3LYP functionals within Reference–State One–Particle Density–Matrix Theory

James P. Finley

Department of Physical Sciences, Eastern New Mexico University,

Station #33, Portales, NM 88130 and*

Department of Applied Chemistry, Graduate School of Engineering,

The University of Tokyo, Tokyo, Japan 113-8656

(Dated: February 2, 2008)

Abstract

For closed-shell systems, the local density approximation (LDA) and the LYP, BLYP, and B3LYP functionals are shown to be compatible with reference-state one-particle density-matrix theory, where this recently introduced formalism is based on Brueckner-orbital theory and an energy functional that includes exact exchange and a non-universal correlation-energy functional. The method is demonstrated to reduce to a density functional theory when the exchange-correlation energy-functional has a simplified form, i.e., its integrand contains only the coordinates of two electron, say $\mathbf{r}_1$ and $\mathbf{r}_2$, and it has a Dirac delta function $\delta(\mathbf{r}_1 - \mathbf{r}_2)$ as a factor. Since Brueckner and Hartree–Fock orbitals are often very similar, any local exchange functional that works well with Hartree–Fock theory is a reasonable approximation with reference-state one-particle density-matrix theory. The LDA approximation is also a reasonable approximation. However, the Colle–Salvetti correlation-energy functional, and the LYP variant, are not ideal for the method, since these are universal functionals. Nevertheless, they appear to provide reasonable approximations. The B3LYP functional is derived using a linear combination of two functionals: One is the BLYP functional; the other uses exact exchange and a correlation-energy functional from the LDA.

PACS numbers:

*Electronic address: james.finley@enmu.edu

I. INTRODUCTION

The Kohn-Sham version of density functional theory (DFT) plays a major role in both quantum chemistry and condensed matter physics [1, 2, 3, 4, 5, 6, 7]. The local density approximation (LDA) [8] has been widely used for the solid state, while for molecules, the most successful functional, a hybrid one [9, 10, 11, 12], is known as B3LYP [9, 13], where this functional contains three parameters, two correlation-energy functionals, the Dirac-exchange functional with the a correction, and exact exchange. The LYP correlation-energy functional [14] – a key component of B3LYP – is derived from the Colle–Salvetti correlation-energy functional [15].

Recently, a density-matrix formalism was developed called reference-state one-particle density-matrix theory [16, 17]. This method employs generalized Hartree–Fock equations containing the exact exchange-potential and a nonlocal correlation-potential, where these equations are obtained using the Brillouin–Brueckner condition (for non-variational energy functionals) or functional minimization (for variational functionals). The method uses energy functionals that yield the target-state energy when the reference state – or its corresponding one-particle density-matrix – is constructed from Brueckner orbitals. Unlike traditional DFT approaches, the v -representable problem does *not* appear in the method, nor the need to introduce functionals defined by a constrained search. The correlation-energy functionals in the method are non-universal, in the sense that they depend on the external potential. Nevertheless, model systems can still be used to derive universal energy-functionals. For example, the Colle–Salvetti functional was shown to be compatible with the method as well as a one-particle-density-matrix variant of the LDA. In addition, using time-independent many-body perturbation theory [18, 19], diagrammatic expansions were given for the non-variational correlation-energy functionals that are expressed in terms of orbitals and orbital energies. When restrictions are placed on the orbital energies, the individual diagrams are shown to *explicitly* depend on the one-particle density-matrix of the reference state (and the external potential).

Below we demonstrate that for systems that can be treated with a simplified exchange-correlation energy-functional – leading to a local exchange-correlation potential – the reference-state one-particle density-matrix formalism reduces to a density functional formalism, where the density that appears within the method is the one from the reference

state, and *not* the target state. This approach differs in this way and in others from the formalism by Lindgren and Salomonson [20], where their method is a generalization of the Hartree–Fock Kohn–Sham approach [21] and yields orbitals that are believed to be very similar to Brueckner orbitals.

The DFT functionals derived from a uniform electron gas are also shown to be compatible with the reference-state one-particle density-matrix theory, where this compatibility appears because the density of the target state for a uniform electron gas is the same as the density of the reference state and, also, the plane-wave states are Brueckner orbitals. In addition, functionals that approximate the exact exchange-energy can also be used, since the exact exchange-functional for Brueckner orbital theory, Hartree–Fock theory, and DFT are the same. Furthermore, functionals derived using the one-particle density-matrix of the reference state are compatible, assuming the errors due to the external potential are small – if the functional is a universal one. Using these observations, the LDA, and the BLYP and B3LYP functionals are shown to be compatible with the reference-state one-particle density-matrix formalism. This puts these functionals within an alternative DFT approach that avoids, for the most part, existence theorems and constraint searches.

II. REFERENCE-STATE ONE-PARTICLE DENSITY-MATRIX THEORY

We seek an eigenfunction $|\Psi\rangle$ on the time-independent Hamiltonian operator:

$$H = \sum_{ij} ([i|(-\frac{1}{2}\nabla^2)|j] + [i|v|j]) a_i^\dagger a_j + \frac{1}{2} \sum_{ijkl} [ij|kl] a_i^\dagger a_k^\dagger a_l a_j, \quad (1)$$

where the integrals are written using chemist’s notation [22]:

$$[ij|kl] = \sum_{\omega_1\omega_2} \int \psi_i^*(\mathbf{x}_1) \psi_j(\mathbf{x}_1) r_{12}^{-1} \psi_k^*(\mathbf{x}_2) \psi_l(\mathbf{x}_2) d\mathbf{r}_1 d\mathbf{r}_2, \quad (2)$$

and the spatial and spin coordinates, $\mathbf{r}$ and ω , are denoted collectively by $\mathbf{x}$. The eigenfunction of interest, or target state $|\Psi\rangle$, yields the electronic energy $\mathcal{E}$:

$$\mathcal{E} = \frac{\langle\Psi|H|\Psi\rangle}{\langle\Psi|\Psi\rangle}. \quad (3)$$

Consider a single-determinantal reference-state, say $|\Phi\rangle$, where this state is a first-order approximation of the target state; the first-order energy is

$$E_1[\Phi] = \langle\Phi|H|\Phi\rangle. \quad (4)$$

Adding and subtracting $E_1[\Phi]$ on the left side of Eq. (3), gives

$$\mathcal{E} = E_1 + (\mathcal{E} - E_1) = E_1[\Phi] + \mathcal{E}_{\text{co}}[\Phi], \quad (5)$$

where this Eq. defines the correlation energy $\mathcal{E}_{\text{co}}[\Phi]$ as $(\mathcal{E} - E_1)$.

There is a one-to-one correspondence between determinant states and their one-particle density-matrices [2, 23], where these density-matrices γ are given by [24, 25, 26, 27, 28]

$$\gamma(\mathbf{x}, \mathbf{x}') = \sum_{w \in \{\psi_o \rightarrow \Phi\}} \psi_w(\mathbf{x}) \psi_w^*(\mathbf{x}'); \quad \gamma \longleftrightarrow |\Phi\rangle, \quad (6)$$

and the notation on the right-hand-side indicates the one-to-one correspondence; furthermore, the sum over w includes only the *occupied* orbitals of the reference state $|\Phi\rangle$: This orbital set is denoted by $\{\psi_o \rightarrow \Phi\}$; the *unoccupied* set is denoted by $\{\psi_u \rightarrow \Phi\}$. Unless stated otherwise, two sets of occupied or unoccupied orbitals that differ by a unitary transformation are considered equivalent.

Because of the one-to-one correspondence mentioned above, any functional that depends on the reference state $|\Phi\rangle$ can be transformed into one that depends on the one-particle density-matrix γ . For example, by requiring the external potential $v(\mathbf{r})$ to be a spin-free operator, the first-order energy can be written as

$$E_1[\gamma] = \int \left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 \gamma(\mathbf{x}, \mathbf{x}') \right]_{\mathbf{x}'=\mathbf{x}} d\mathbf{x} + \int v(\mathbf{r}) \gamma(\mathbf{x}, \mathbf{x}) d\mathbf{x} + E_J[\gamma] + E_x[\gamma], \quad (7)$$

where the Coulomb and exchange energies have their usual forms:

$$E_J[\gamma] = \frac{1}{2} \iint r_{12}^{-1} \gamma(\mathbf{x}_1, \mathbf{x}_1) \gamma(\mathbf{x}_2, \mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2, \quad (8)$$

$$-E_x[\gamma] = \frac{1}{2} \iint r_{12}^{-1} \gamma(\mathbf{x}_1, \mathbf{x}_2) \gamma(\mathbf{x}_2, \mathbf{x}_1) d\mathbf{x}_1 d\mathbf{x}_2, \quad (9)$$

and, henceforth, an integration over $\mathbf{x}$ implies an integration over $\mathbf{r}$ and a summation over ω .

A. Hartree–Fock Theory

Let the Fock operator be denoted $\hat{F}_\gamma$; the functional derivative of the first-order energy $E_1[\gamma]$ gives the kernel of this operator [2]:

$$\begin{aligned} F_\gamma(\mathbf{x}_1, \mathbf{x}_2) &= \frac{\delta E_1[\gamma]}{\delta \gamma(\mathbf{x}_2, \mathbf{x}_1)} \\ &= \delta(\mathbf{x}_2 - \mathbf{x}_1) \left(-\frac{1}{2} \nabla_2^2 + v(\mathbf{r}_2) + \int r_{23}^{-1} \gamma(\mathbf{x}_3, \mathbf{x}_3) d\mathbf{x}_3 \right) + v_x^\gamma(\mathbf{x}_1, \mathbf{x}_2), \end{aligned} \quad (10)$$

where the two-body function, $v_x^\gamma(\mathbf{x}_1, \mathbf{x}_2)$, is obtained from the exchange energy:

$$v_x^\gamma(\mathbf{x}_1, \mathbf{x}_2) = \frac{\delta E_x[\gamma]}{\delta \gamma(\mathbf{x}_2, \mathbf{x}_1)} = -r_{12}^{-1} \gamma(\mathbf{x}_1, \mathbf{x}_2), \quad (11)$$

and this function is the kernel of the exchange operator, denoted by $\hat{v}_x^\gamma$.

Minimizing the functional $E_1[\gamma]$, subject to the constraint that the one-particle density-matrix comes from a single-determinantal state, yields $E_1[\tilde{\tau}]$, where $\tilde{\tau}$ is the one-particle density-matrix of the Hartree–Fock reference-state $|\tilde{\Theta}\rangle$:

$$\tilde{\tau}(\mathbf{x}, \mathbf{x}') = \sum_{w \in \{\psi_o \rightarrow \tilde{\Theta}\}} \psi_w(\mathbf{x}) \psi_w^*(\mathbf{x}'), \quad (12)$$

and this density-matrix satisfies the following condition:

$$\langle \psi_r | \hat{F}_{\tilde{\tau}} | \psi_w \rangle = 0; \quad \psi_w \in \{\psi_o \rightarrow \tilde{\tau}\}, \quad \psi_r \in \{\psi_u \rightarrow \tilde{\tau}\}, \quad (13a)$$

and the right-hand-side notation indicates that ψ_w is a member of the set of occupied orbital that determines $\tilde{\tau}$ (and $|\tilde{\Theta}\rangle$); ψ_r is unoccupied. The operator form of this Eq. is

$$(\hat{1} - \tilde{\tau}) \hat{F}_{\tilde{\tau}} \tilde{\tau} = 0, \quad (13b)$$

where the identity operator $\hat{1}$, using a Hartree–Fock basis set, is given by

$$\hat{1} = \sum_{w \in \{\psi_o \rightarrow \tilde{\tau}\}} |\psi_w\rangle \langle \psi_w| + \sum_{r \in \{\psi_u \rightarrow \tilde{\tau}\}} |\psi_r\rangle \langle \psi_r|. \quad (14)$$

A unique set of occupied and unoccupied orbitals is obtained by requiring the occupied and unoccupied blocks of $\hat{F}_{\tilde{\tau}}$ to be diagonal:

$$\hat{F}_{\tilde{\tau}} \psi_w^{\tilde{\tau}}(\mathbf{x}) = \varepsilon_w^{\tilde{\tau}} \psi_w^{\tilde{\tau}}(\mathbf{x}), \quad \psi_w^{\tilde{\tau}} \in \{\psi_o \rightarrow \tilde{\tau}\}, \quad (15a)$$

$$\hat{F}_{\tilde{\tau}} \psi_r^{\tilde{\tau}}(\mathbf{x}) = \varepsilon_r^{\tilde{\tau}} \psi_r^{\tilde{\tau}}(\mathbf{x}), \quad \psi_r^{\tilde{\tau}} \in \{\psi_u \rightarrow \tilde{\tau}\}. \quad (15b)$$

The orbitals sets that satisfy Eqs. (13) and (15) are denoted by $\{\psi_o^{\tilde{\tau}} \leftarrow \tilde{\tau}, \hat{F}_{\tilde{\tau}}\}$ and $\{\psi_u^{\tilde{\tau}} \leftarrow \tilde{\tau}, \hat{F}_{\tilde{\tau}}\}$, indicating that they are uniquely determined by $\tilde{\tau}$ and $\hat{F}_{\tilde{\tau}}$.

Substituting Eq. (10) into Eqs. (15) gives the canonical Hartree–Fock Eqs:

$$\left(-\frac{1}{2} \nabla_1^2 + v(\mathbf{r}_1) + \int r_{12}^{-1} \tilde{\tau}(\mathbf{x}_2, \mathbf{x}_2) d\mathbf{x}_2 + \hat{v}_x^{\tilde{\tau}}(\mathbf{x}_1) \right) \psi_i^{\tilde{\tau}}(\mathbf{x}_1) = \varepsilon_i^{\tilde{\tau}} \psi_i^{\tilde{\tau}}(\mathbf{x}_1). \quad (16)$$

B. Variational Brueckner orbital theory

Brueckner orbital theory [20, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43] is a generalization of Hartree–Fock theory that utilizes a single-determinantal state that has the maximum overlap with the target state [44, 45]. We now review a variant of Brueckner orbital theory that is used to derive the reference-state one-particle density-matrix theory [17].

In this previous work, we introduced four trial wavefunctions – say $|\Psi_{\Phi}^{(\eta)}\rangle$, where ($\eta =$ I, II, III, and IV) – that are defined with respect to a single-determinantal reference-state $|\Phi\rangle$. The first trial-wavefunction $|\Psi_{\Phi}^{(I)}\rangle$ is simply the target state of interest, $|\Psi\rangle$, with the single excitations removed:

$$|\Psi_{\Phi}^{(I)}\rangle = (1 - P_{11}^{\Phi}) |\Psi\rangle, \quad (17)$$

where the projector for the singly-excited states is

$$P_{11}^{\Phi} = \sum_{w \in \{\psi_o \rightarrow \Phi\}} \sum_{r \in \{\psi_u \rightarrow \Phi\}} |\Phi_w^r\rangle \langle \Phi_w^r|, \quad (18)$$

and the singly-excited state are

$$|\Phi_w^r\rangle = a_r^{\dagger} a_w |\Phi\rangle; \quad \psi_w \in \{\psi_o \rightarrow \Phi\}, \quad \psi_r \in \{\psi_u \rightarrow \Phi\}. \quad (19)$$

The P_{11}^{Φ} subspace is completely determined by $|\Phi\rangle$; P_{11}^{Φ} is also invariant to a unitary transformation of occupied, or virtual, orbitals [36].

The second trial-wavefunction $|\Psi_{\Phi}^{(II)}\rangle$ is defined with respect to the target state expressed by an exponential ansatz: ($|\Psi\rangle = e^{S_{\Phi}} |\Phi\rangle$), where $|\Psi_{\Phi}^{(II)}\rangle$ is generated by removing the single-excitation amplitudes S_1^{Φ} from the cluster-operator S :

$$|\Psi_{\Phi}^{(II)}\rangle = e^{(S_{\Phi} - S_1^{\Phi})} |\Phi\rangle. \quad (20)$$

The third trial-wavefunction $|\Psi_{\Phi}^{(III)}\rangle$ can be generated by its wave-operator:

$$\hat{\Omega}_{\Phi} |\Phi\rangle = |\Psi_{\Phi}^{(III)}\rangle, \quad (21)$$

that can be expressed in an exponential form: ($\hat{\Omega}_{\Phi} = e^{\hat{S}_{\Phi}} |\Phi\rangle$), where $\hat{S}_{\Phi}$ can be written as a sum n -body excitations, with the exclusion of a one-body operator:

$$\hat{S}_{\Phi} = \hat{S}_2^{\Phi} + \hat{S}_3^{\Phi} + \dots. \quad (22)$$

The wave operator $\hat{\Omega}_\Phi$ is as a solution to the coupled cluster equations [18, 19, 46, 47, 48, 49, 50, 51, 52, 53] with the single excitation portion removed:

$$(1 - P_{11}^\Phi) \left(H \hat{\Omega}_\Phi \right)_{\text{op,cn}} = 0, \quad (23)$$

where only the open (op) and connected (cn) portions enter into the relation. This expression defines the trial functional $|\Psi_\Phi^{(\text{III})}\rangle$ using Eq. (21).

The fourth trial wavefunctions $|\Psi_\Phi^{(\text{IV})}\rangle$ is not considered here, except to mention that it is obtained by solving the configuration-interaction equations [19, 22, 54] in an approximate way, i.e., by neglecting the single-excitation portion.

All of the trial states $|\Psi_\Phi^{(\eta)}\rangle$ share the property that they contain no single excitations. i.e., $(P_{11}^\Phi |\Psi_\Phi^{(\eta)}\rangle = 0)$, and they generate the target state $|\Psi\rangle$ when their reference state satisfies $(|\Phi\rangle = |\Theta\rangle)$, where $|\Theta\rangle$ is the determinantal state constructed from occupied Bruckner orbitals. In other words, we have

$$|\Psi_\Theta^{(\eta)}\rangle = |\Psi\rangle. \quad (24)$$

Because of the one-to-one correspondence mentioned above, the trial wavefunctions depend on the one-particle density-matrix; so we can write $|\Psi_\gamma^{(\eta)}\rangle$. Using this notation, we can define variational energy-functionals that depend on the one-particle density-matrix:

$$\bar{E}_\eta[\gamma] = \frac{\langle \Psi_\gamma^{(\eta)} | H | \Psi_\gamma^{(\eta)} \rangle}{\langle \Psi_\gamma^{(\eta)} | \Psi_\gamma^{(\eta)} \rangle} = E_1[\gamma] + \bar{E}_{\text{co}}^{(\eta)}[\gamma], \quad (25)$$

where the last relation defines the correlation-energy functionals $\bar{E}_{\text{co}}^{(\eta)}[\gamma]$ as $(\bar{E}_\eta[\gamma] - E_1[\gamma])$.

Substituting Eq. (7) into Eqs. (5) and (25) gives the following:

$$\mathcal{E} = \int \left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 \gamma(\mathbf{x}, \mathbf{x}') \right]_{\mathbf{x}'=\mathbf{x}} d\mathbf{x} + \int v(\mathbf{r}) \gamma(\mathbf{x}, \mathbf{x}) d\mathbf{x} + E_J[\gamma] + \mathcal{E}_{\text{xc}}[\gamma], \quad (26a)$$

$$\bar{E}_\eta[\gamma] = \int \left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 \gamma(\mathbf{x}, \mathbf{x}') \right]_{\mathbf{x}'=\mathbf{x}} d\mathbf{x} + \int v(\mathbf{r}) \gamma(\mathbf{x}, \mathbf{x}) d\mathbf{x} + E_J[\gamma] + \bar{E}_{\text{xc}}^{(\eta)}[\gamma], \quad (26b)$$

where the exchange-correlation energy and exchange-correlation energy-functionals are, respectively, defined by

$$\mathcal{E}_{\text{xc}}[\gamma] = E_{\text{x}}[\gamma] + \mathcal{E}_{\text{co}}[\gamma], \quad (27a)$$

$$\bar{E}_{\text{xc}}^{(\eta)}[\gamma] = E_{\text{x}}[\gamma] + \bar{E}_{\text{co}}^{(\eta)}[\gamma]. \quad (27b)$$

We now generalize many of the Hartree–Fock relations from the previous subsection simply by replacing $E_1[\gamma]$ with $\bar{E}_\eta[\gamma]$ and the Fock operator $\hat{F}_\gamma$ with generalized, or exact, Fock operators, $\hat{\zeta}_\gamma^{(\eta)}$.

The functional derivative of our energy functionals yield two-body functions that serve as the kernels of the exact Fock operators:

$$\begin{aligned}\zeta_\gamma^{(\eta)}(\mathbf{x}_1, \mathbf{x}_2) &= \frac{\bar{E}_\eta[\gamma]}{\delta\gamma(\mathbf{x}_2, \mathbf{x}_1)} \\ &= \delta(\mathbf{x}_2 - \mathbf{x}_1) \left(-\frac{1}{2}\nabla_2^2 + v(\mathbf{r}_2) + \int r_{23}^{-1} \gamma(\mathbf{x}_3, \mathbf{x}_3) d\mathbf{x}_3 \right) + \nu_{\text{xc}}^{\gamma\eta}(\mathbf{x}_1, \mathbf{x}_2),\end{aligned}\tag{28}$$

where the kernels of the exchange-correlation operators, $\nu_{\text{xc}}^{\gamma\eta}(\mathbf{x}_1, \mathbf{x}_2)$, are obtained from the exchange-correlation energy-functionals:

$$\nu_{\text{xc}}^{\gamma\eta}(\mathbf{x}_1, \mathbf{x}_2) = \frac{\delta\bar{E}_{\text{xc}}^{(\eta)}[\gamma]}{\delta\gamma(\mathbf{x}_2, \mathbf{x}_1)} = \frac{\delta\bar{E}_{\text{co}}^{(\eta)}[\gamma]}{\delta\gamma(\mathbf{x}_2, \mathbf{x}_1)} - r_{12}^{-1}\gamma(\mathbf{x}_1, \mathbf{x}_2),\tag{29}$$

where the last relation uses Eqs. (27b) and (11).

We now require the target state to be a ground state; using the variation theorem and Eq. (24), the minimizing of the functionals $\bar{E}_\eta[\gamma]$ – subject to the constraint that the one-particle density-matrix comes from a single-determinantal state – yields

$$\mathcal{E} = \bar{E}_\eta[\tau],\tag{30}$$

and from Eqs. (26) and (27) we have

$$\mathcal{E}_{\text{co}}[\tau] = \bar{E}_{\text{co}}^{(\eta)}[\tau],\tag{31}$$

$$\mathcal{E}_{\text{xc}}[\tau] = \bar{E}_{\text{xc}}^{(\eta)}[\tau],\tag{32}$$

where τ is the one-particle density-matrix of the Brueckner reference-state $|\Theta\rangle$:

$$\tau(\mathbf{x}, \mathbf{x}') = \sum_{w \in \{\psi_o \rightarrow \Theta\}} \psi_w(\mathbf{x}) \psi_w^*(\mathbf{x}'),\tag{33}$$

and the Brueckner orbitals satisfy the following equivalent conditions:

$$\langle \psi_r | \hat{\zeta}_\tau^{(\eta)} | \psi_w \rangle = 0; \quad \psi_w \in \{\psi_o \rightarrow \tau\}, \quad \psi_r \in \{\psi_u \rightarrow \tau\},\tag{34a}$$

$$(\hat{1} - \tau) \hat{\zeta}_\tau^{(\eta)} \tau = 0,\tag{34b}$$

where these orbitals do not depend of η – any trial wavefunction gives the same results.

A unique set of occupied and unoccupied orbitals is obtained by requiring the occupied and unoccupied blocks of $\hat{\zeta}_\tau^{(\eta)}$ to be diagonal:

$$\hat{\zeta}_\tau^{(\eta)} \psi_w^\tau(\mathbf{x}) = \xi_w^\tau \psi_w^\tau(\mathbf{x}), \quad \psi_w^\tau \in \{\psi_o \rightarrow \tau\},\tag{35a}$$

$$\hat{\zeta}_\tau^{(\eta)} \psi_r^\tau(\mathbf{x}) = \xi_r^\tau \psi_r^\tau(\mathbf{x}), \quad \psi_r^\tau \in \{\psi_u \rightarrow \tau\}.\tag{35b}$$

Henceforth, the orbitals sets that satisfy Eqs. (34) and (35) are denoted by $\{\psi_o^\tau \leftarrow \tau, \hat{\zeta}_\tau^{(\eta)}\}$ and $\{\psi_u^\tau \leftarrow \tau, \hat{\zeta}_\tau^{(\eta)}\}$, indicating that they are determined by τ and $\hat{\zeta}_\tau^{(\eta)}$. Since theses orbitals, and their energies, can, perhaps, depend on η , it is more precise to denote then by $\psi_i^{\tau\eta}$ and $\xi_i^{\tau\eta}$, but we suppress the η superscripts to keep the notation less cluttered.

Substituting Eq. (28) into Eqs. (35) gives generalized, canonical Hartree–Fock Eqs:

$$\left(-\frac{1}{2}\nabla_1^2 + v(\mathbf{r}_1) + \int r_{12}^{-1}\tau(\mathbf{x}_2, \mathbf{x}_2) d\mathbf{x}_2 + \hat{\nu}_{xc}^{\tau\eta}(\mathbf{x}_1) \right) \psi_i^\tau(\mathbf{x}_1) = \xi_i^\tau \psi_i^\tau(\mathbf{x}_1). \quad (36)$$

III. CLOSED-SHELL CASE

Our interest here is in cases where the Hamiltonian is spin-free – it contains no spin coordinates – and the target state $|\Psi\rangle$ is a ground-state singlet that is well described by a closed-shell reference-state $|\Phi\rangle$. In these cases – and many others – it is appropriate to use a set of spatially restricted orbitals, given by

$$\psi_{j\sigma}(\mathbf{x}) = \chi_j(\mathbf{r})\sigma(\omega); \quad \sigma = \alpha, \beta, \quad (37)$$

so that $|\Phi\rangle$ is determined by a set of doubly-occupied spatial orbitals, denoted by $\{\chi_o \rightarrow \Phi\}$, where this set also determines the virtual set, denoted by $\{\chi_u \rightarrow \Phi\}$. Two sets of orbitals that differ by a unitary transformation are, again, considered equivalent.

By definition, the spinless, one-particle density-matrix is given by [2, 28]

$$\rho_1(\mathbf{r}_1, \mathbf{r}_2) = \sum_{\omega} \gamma(\mathbf{r}_1, \omega; \mathbf{r}_2, \omega), \quad (38)$$

and from Eqs. (37) and (6) with ($w = x\sigma$) we have

$$\rho_1(\mathbf{r}_1, \mathbf{r}_2) = 2 \sum_{x \in \{\chi_o \rightarrow \Phi\}} \chi_x(\mathbf{r}_1) \chi_x^*(\mathbf{r}_2); \quad \rho_1 \longleftrightarrow |\Phi\rangle. \quad (39)$$

By using the restricted orbitals defined by Eq. (37), it is easily demonstrated that the one-particle density-matrix γ is determined by the spinless one, as indicated by the following relation:

$$\gamma(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{2} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \delta_{\omega_1 \omega_2}. \quad (40)$$

Hence, any functional of γ now becomes a functional of ρ_1 . In particular, for a spin-free Hamiltonian, the spin-variable summations are easily performed, yielding explicit functionals

of ρ_1 . For example, the electronic energy and our energy-functionals, Eqs. (5) and (25), can be written as

$$\mathcal{E} = E_1[\rho_1] + \mathcal{E}_{\text{co}}[\rho_1], \quad (41a)$$

$$\bar{E}_\eta[\rho_1] = E_1[\rho_1] + \bar{E}_{\text{co}}^{(\eta)}[\rho_1], \quad (41b)$$

or the representations of these functionals, given by Eqs. (26), reduce to

$$\mathcal{E} = \int \left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 \rho_1(\mathbf{r}, \mathbf{r}') \right]_{\mathbf{r}'=\mathbf{r}} d\mathbf{r} + \int v(\mathbf{r}) \rho(\mathbf{r}) d\mathbf{r} + E_{\text{J}}[\rho] + \mathcal{E}_{\text{xc}}[\rho_1], \quad (42a)$$

$$\bar{E}_\eta[\rho_1] = \int \left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 \rho_1(\mathbf{r}, \mathbf{r}') \right]_{\mathbf{r}'=\mathbf{r}} d\mathbf{r} + \int v(\mathbf{r}) \rho(\mathbf{r}) d\mathbf{r} + E_{\text{J}}[\rho] + \bar{E}_{\text{xc}}^{(\eta)}[\rho_1], \quad (42b)$$

where the definitions given by Eqs. (27) can be expressed using a dependence upon ρ_1 :

$$\mathcal{E}_{\text{xc}}[\rho_1] = E_{\text{x}}[\rho_1] + \mathcal{E}_{\text{co}}[\rho_1], \quad (43a)$$

$$\bar{E}_{\text{xc}}^{(\eta)}[\rho_1] = E_{\text{x}}[\rho_1] + \bar{E}_{\text{co}}^{(\eta)}[\rho_1]; \quad (43b)$$

furthermore, the first-order energy, Eq. (7), becomes

$$E_1[\rho_1] = \int \left[-\frac{1}{2} \nabla_{\mathbf{r}}^2 \rho_1(\mathbf{r}, \mathbf{r}') \right]_{\mathbf{r}'=\mathbf{r}} d\mathbf{r} + \int v(\mathbf{r}) \rho(\mathbf{r}) d\mathbf{r} + E_{\text{J}}[\rho] + E_{\text{x}}[\rho_1], \quad (44)$$

where the Coulomb and exchange energies, Eq. (8) and (9), reduce to

$$E_{\text{J}}[\rho] = \frac{1}{2} \iint r_{12}^{-1} \rho(\mathbf{r}_1) \rho(\mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2, \quad (45)$$

$$-E_{\text{x}}[\rho_1] = \frac{1}{4} \iint r_{12}^{-1} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1) d\mathbf{r}_1 d\mathbf{r}_2, \quad (46)$$

and the particle density is

$$\rho(\mathbf{r}) = \rho_1(\mathbf{r}, \mathbf{r}). \quad (47)$$

A. Hartree–Fock Theory

Using Eqs. (11) and (40) it is readily observed that the kernel of the exchange operator is given by

$$v_{\text{x}}^{\gamma}(\mathbf{x}_1, \mathbf{x}_2) = v_{\text{x}}^{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) \delta_{\omega_1, \omega_2}, \quad (48)$$

where

$$v_{\text{x}}^{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) = -\frac{1}{2} r_{12}^{-1} \rho_1(\mathbf{r}_1, \mathbf{r}_2). \quad (49)$$

Therefore, for an arbitrary function, say ϕ , we have

$$\hat{v}_x^\gamma \phi(\mathbf{x}_1) = \hat{v}_x^{\rho_1} \phi(\mathbf{x}_1), \quad (50)$$

where $v_x^\gamma(\mathbf{x}_1, \mathbf{x}_2)$ and $v_x^{\rho_1}(\mathbf{r}_1, \mathbf{r}_2)$ serve as the kernels of $\hat{v}_x^\gamma$ and $\hat{v}_x^{\rho_1}$, respectively:

$$\hat{v}_x^\gamma \phi(\mathbf{x}_1) = \int v_x^\gamma(\mathbf{x}_1, \mathbf{x}_2) \phi(\mathbf{x}_2) d\mathbf{x}_2, \quad (51)$$

$$\hat{v}_x^{\rho_1} \phi'(\mathbf{r}_1) = \int v_x^{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) \phi'(\mathbf{r}_2) d\mathbf{r}_2. \quad (52)$$

Using Eqs. (48) and (40), the kernel of the Fock operators, Eq. (10), can be written as

$$F_\gamma(\mathbf{x}_1, \mathbf{x}_2) = \delta_{\omega_1 \omega_2} F_{\rho_1}(\mathbf{r}_1, \mathbf{r}_2), \quad (53a)$$

where

$$F_{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) = \delta(\mathbf{r}_2 - \mathbf{r}_1) \left(-\frac{1}{2} \nabla_2^2 + v(\mathbf{r}_2) + \int r_{23}^{-1} \rho(\mathbf{r}_3) d\mathbf{r}_3 \right) + v_x^{\rho_1}(\mathbf{r}_1, \mathbf{r}_2), \quad (53b)$$

Substituting Eqs. (37) and (53a) into (15), and summing over ω_2 we get

$$\int F_{\tilde{\rho}_1}(\mathbf{r}_1, \mathbf{r}_2) \chi_j^{\tilde{\rho}_1}(\mathbf{r}_2) d\mathbf{r}_2 \sigma(\omega_1) = \varepsilon_j^{\tilde{\rho}_1} \chi_j^{\tilde{\rho}_1}(\mathbf{r}_1) \sigma(\omega_1). \quad (54)$$

By letting $F_{\tilde{\rho}_1}(\mathbf{r}_1, \mathbf{r}_2)$ serve as the kernel of the spin-free Fock-operator $\hat{F}_{\tilde{\rho}_1}$, we have

$$\hat{F}_{\tilde{\rho}_1} \chi_j^{\tilde{\rho}_1}(\mathbf{r}_1) = \varepsilon_j^{\tilde{\rho}_1} \chi_j^{\tilde{\rho}_1}(\mathbf{r}_1), \quad (55)$$

where $\tilde{\rho}_1$ is the spinless one-particle density-matrix of the Hartree-Fock state:

$$\tilde{\tau}(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{2} \tilde{\rho}_1(\mathbf{r}_1, \mathbf{r}_2) \delta_{\omega_1 \omega_2}, \quad (56)$$

and the Hartree-Fock spatial-orbitals are denoted in an analogous way as the spin-orbitals from Sec. (II A), e.g., $\tilde{\rho}_1$ replaces $\tilde{\tau}$.

Substituting Eqs. (53b) into (54) and using (52) gives

$$\left(-\frac{1}{2} \nabla_1^2 + v(\mathbf{r}_1) + \int r_{12}^{-1} \tilde{\rho}(\mathbf{r}_2) d\mathbf{r}_2 - \hat{v}_x^{\tilde{\rho}_1}(\mathbf{r}_1) \right) \chi_i^{\tilde{\rho}_1}(\mathbf{r}_1) = \varepsilon_i^{\tilde{\rho}_1} \chi_i^{\tilde{\rho}_1}(\mathbf{r}_1). \quad (57)$$

Eqs. (55) and (57) are the closed-shell spin-free forms of Eqs. (15) and (16); the spin-free forms of Eqs. (10) and (11) are easily proven to satisfy the following relations:

$$F_{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) = \frac{\delta E_1[\rho_1]}{\delta \rho_1(\mathbf{r}_2, \mathbf{r}_1)}, \quad (58)$$

$$v_x^{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) = \frac{\delta E_x[\rho_1]}{\delta \rho_1(\mathbf{r}_2, \mathbf{r}_1)}, \quad (59)$$

where, as mentioned previously, these functions are given by Eqs. (53b) and (49), respectively.

B. Brueckner orbital theory

We now generalize many of the Hartree–Fock relations from the previous subsection simply by replacing the exchange operator $\hat{v}_x^\gamma$ with the exchange-correlation operator $\hat{\nu}_{xc}^{\gamma\eta}$.

For the closed shell case under consideration, it can be easily demonstrated that the kernels of the exchange-correlation operators are given by [55]

$$\nu_{xc}^{\gamma\eta}(\mathbf{x}_1, \mathbf{x}_2) = \nu_{xc}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2)\delta_{\omega_1, \omega_2}. \quad (60)$$

Therefore, for an arbitrary function, say ϕ , we have

$$\hat{\nu}_{xc}^{\gamma\eta}\phi(\mathbf{x}_1) = \hat{\nu}_{xc}^{\rho_1\eta}\phi(\mathbf{x}_1), \quad (61)$$

where $\nu_{xc}^{\gamma\eta}(\mathbf{x}_1, \mathbf{x}_2)$ and $\nu_{xc}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2)$ serve as the kernels of $\hat{\nu}_{xc}^{\gamma\eta}$ and $\hat{\nu}_{xc}^{\rho_1\eta}$, respectively.

Using Eqs. (60) and (40), the kernel of the exact Fock operators, Eq. (28), can be written as

$$\zeta_\gamma^{(\eta)}(\mathbf{x}_1, \mathbf{x}_2) = \delta_{\omega_1\omega_2}\zeta_{\rho_1}^{(\eta)}(\mathbf{r}_1, \mathbf{r}_2), \quad (62a)$$

where

$$\zeta_{\rho_1}^{(\eta)}(\mathbf{r}_1, \mathbf{r}_2) = \delta(\mathbf{r}_2 - \mathbf{r}_1) \left(-\frac{1}{2}\nabla_2^2 + v(\mathbf{r}_2) + \int r_{23}^{-1}\rho(\mathbf{r}_3) d\mathbf{r}_3 \right) + \nu_{xc}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2). \quad (62b)$$

Substituting Eqs. (37) and (62a) into (35), and summing over ω_2 we get

$$\int \zeta_{\rho_1}^{(\eta)}(\mathbf{r}_1, \mathbf{r}_2) \chi_j^{\rho_1}(\mathbf{r}_2) d\mathbf{r}_2 \sigma(\omega_1) = \xi_j^{\rho_1} \chi_j^{\rho_1}(\mathbf{r}_1) \sigma(\omega_1). \quad (63)$$

By letting $\zeta_{\rho_1}^{(\eta)}(\mathbf{r}_1, \mathbf{r}_2)$ serve as the kernels of the exact, spin-free Fock-operator $\hat{\zeta}_{\rho_1}^{(\eta)}$, we have

$$\hat{\zeta}_{\rho_1}^{(\eta)} \chi_j^{\rho_1}(\mathbf{r}_1) = \xi_j^{\rho_1} \chi_j^{\rho_1}(\mathbf{r}_1), \quad (64)$$

where ρ_1 is the spinless one-particle density of the Brueckner state:

$$\tau(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{2} \rho_1(\mathbf{r}_1, \mathbf{r}_2) \delta_{\omega_1\omega_2}, \quad (65)$$

and the Brueckner spatial-orbitals are denoted in an analogous way as the spin-orbitals from Sec. (II B), e.g., ρ_1 replaces τ .

Substituting Eqs. (62b) into (63) gives

$$\left(-\frac{1}{2}\nabla_1^2 + v(\mathbf{r}_1) + \int r_{12}^{-1}\rho(\mathbf{r}_2) d\mathbf{r}_2 + \hat{\nu}_{xc}^{\rho_1\eta}(\mathbf{r}_1) \right) \chi_i^{\rho_1}(\mathbf{r}_1) = \xi_i^{\rho_1} \chi_i^{\rho_1}(\mathbf{r}_1). \quad (66)$$

Eqs. (64) and (66) are the closed-shell spin-free forms of Eqs. (35) and (36); the spin-free forms or Eqs. (28) and (29) can be shown to satisfy the following relations [55]:

$$\zeta_{\rho_1}^{(\eta)}(\mathbf{r}_1, \mathbf{r}_2) = \frac{\delta \bar{E}_\eta[\rho_1]}{\delta \rho_1(\mathbf{r}_2, \mathbf{r}_1)}, \quad (67)$$

$$\nu_{xc}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2) = \frac{\delta \bar{E}_{xc}^{(\eta)}[\rho_1]}{\delta \rho_1(\mathbf{r}_2, \mathbf{r}_1)}. \quad (68)$$

Substituting Eq. (43b) into (68) and using (59), we have

$$\nu_{xc}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2) = v_x^{\rho_1}(\mathbf{r}_1, \mathbf{r}_2) + \nu_{co}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2), \quad (69)$$

where the kernel of the exchange potential $v_x^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2)$ is given by Eq. (49); the kernel of the correlation potential $\nu_{co}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2)$ is defined by

$$\nu_{co}^{\rho_1\eta}(\mathbf{r}_1, \mathbf{r}_2) = \frac{\delta \bar{E}_{co}^{(\eta)}[\rho_1]}{\delta \rho_1(\mathbf{r}_2, \mathbf{r}_1)}. \quad (70)$$

The operator form of Eq. (69) is given by

$$\hat{\nu}_{xc}^{\rho_1\eta} = \hat{v}_x^{\rho_1} + \hat{\nu}_{co}^{\rho_1\eta}. \quad (71)$$

IV. APPROXIMATIONS

The target state $|\Psi\rangle$ can be generated from a reference state $|\Phi\rangle$ by its wave-operator Ω_Φ :

$$|\Psi\rangle = \Omega_\Phi|\Phi\rangle = |\Psi_\Phi\rangle + |\Psi_{\Phi,\Theta}\rangle, \quad (72)$$

where we have partitioned the target state into two terms: $|\Psi_\Phi\rangle$ and $|\Psi_{\Phi,\Theta}\rangle$. Consider the diagrammatic perturbation-expansion for $\Omega_\Phi|\Phi\rangle$, $|\Psi_{\Phi,\Theta}\rangle$ is then defined as the sum of *all* diagrammatic terms from the perturbation expansion that must vanish for any external potential when the reference state is the Brueckner one:

$$|\Psi_{\Theta,\Theta}\rangle = 0. \quad (73)$$

Hence, $|\Psi_\Phi\rangle$ is a trial wavefunction that fulfills all of the requirements of the ones considered in Sec. II B; furthermore, by our definition, $|\Psi_\Phi\rangle$ is the best trial wavefunction, since it contains no extra terms that must vanish when $(|\Phi\rangle = |\Theta\rangle)$.

Consider a system with an external potential, say v' , where its correlation energy $\mathcal{E}_{co}$ is known when the reference state is the Brueckner one with a corresponding one-particle density-matrix of τ' ; using Eq. (31) we have

$$\mathcal{E}_{co}[\tau', v'] = \bar{E}_{co}[\tau', v'], \quad (74)$$

where the correlation-energy functional from $|\Psi_\Phi\rangle$ is denoted by $\bar{E}_{\text{co}}$ and we have indicated that these functionals depend on the external potential. Hence, a reasonable approximation for $\bar{E}_{\text{co}}$ at other potentials v and one-particle density-matrices γ is given by

$$\bar{E}_{\text{co}}[\gamma, v] = \mathcal{E}_{\text{co}}[\tau', v']_{(\tau'=\gamma, v'=v)}. \quad (75)$$

Now consider the four trial wavefunctions which can also be obtained using a wave operator:

$$|\Psi_\Phi^{(\eta)}\rangle = \Omega_\Phi^{(\eta)}|\Phi\rangle, \quad (76)$$

where from the previous notation we have $(\Omega_\Phi^{(\text{III})} = \tilde{\Omega}_\Phi)$. Consider the first two cases: ($\eta = \text{I}$ and II), where the wave-operators expansion are easily obtained by considering Lindgren's formalism [18, 51, 56]. By examining this expansion, it is readily seen that the trial wavefunctions contain terms, say $|\Psi_{\Phi, \Theta}^{(\eta)}\rangle$, that, by necessity, vanish when $(|\Phi\rangle = |\Theta\rangle)$:

$$|\Psi_\Phi^{(\eta)}\rangle = |\Psi_\Phi^{(\eta)}\rangle + |\Psi_{\Phi, \Theta}^{(\eta)}\rangle. \quad (77)$$

Hence, even though the approximation (75) is reasonable for ($\eta = \text{I}$ and II):

$$\bar{E}_{\text{co}}^{(\eta)}[\gamma, v] = \mathcal{E}_{\text{co}}[\tau', v']_{(\tau'=\gamma, v'=v)}, \quad (78)$$

it neglects the contribution to $\bar{E}_{\text{co}}^{(\eta)}$ arising from $|\Psi_{\Phi, \Theta}^{(\eta)}\rangle$.

The wave operator diagrams for ($\eta = \text{III}$) are more complicated to describe and are given elsewhere [55]. However, it is obvious from the previous work [17] that the term $|\Psi_{\Phi, \Theta}^{(\text{III})}\rangle$ is considerably smaller in magnitude than for ($\eta = \text{I}$ and II), suggesting that the approximation (78) is reasonable, where this type of approximation was considered previously [17]. It is quite possible that $|\Psi_{\Phi, \Theta}^{(\text{III})}\rangle$ is zero, so, in that case ($|\Psi_\Phi^{(\text{III})}\rangle = |\Psi_\Phi\rangle$), this was previously presumed to be true [17], but it was not discussed in detail.

The wave-operator diagrams for ($\eta = \text{IV}$) contain unlinked terms and will not be considered, except to mention that $|\Psi_{\Phi, \Theta}^{(\text{IV})}\rangle$ is nonzero.

A. Local Density Approximation

The correlation energy of an electron gas – denoted by $\mathcal{E}_{\text{co}}^{(\text{gas})}$ – is given by the correlation energy $\mathcal{E}_{\text{co}}$ for a system where the external potential v is constant. Since the orbitals for an

electron gas are Brueckner, and using Eq. (31), we have

$$\mathcal{E}_{\text{co}}^{(\text{gas})}[\tau_g] = \mathcal{E}_{\text{co}}[\tau_g, v] = \bar{E}_{\text{co}}^{(\eta)}[\tau_g, v], \quad v = \text{const}, \quad (79)$$

where τ_g is the one-particle density-matrix that is from the Brueckner reference-state of an electron gas, say $|\Theta_g\rangle$, where the Brueckner orbitals, defining $|\Theta_g\rangle$, are given by plane waves when periodic boundary conditions are imposed; furthermore, the correlation energy $\mathcal{E}_{\text{co}}$ is unchanged if v is changed by a constant amount; $\mathcal{E}_{\text{co}}^{(\text{gas})}[\tau_g]$ does not depend on v . (A constant shift in v does, however, shift the first-order energy E_1 .)

The electron-gas approximation for the third correlation-energy functional utilizes this functional, and it is given by [17]:

$$\bar{E}_{\text{co}}^{(\text{III})}[\gamma] \approx \mathcal{E}_{\text{co}}^{(\text{gas})}[\tau_g]_{(\tau_g=\gamma)}, \quad (80)$$

and this approximation is a special case of Eq. (78), where there are no terms involving an external potential. (We can also use this approximation for $\bar{E}_{\text{co}}$ defined in the previous subsection.)

For spatially restricted orbitals, given by Eq. (37), the above Eq. becomes

$$\bar{E}_{\text{co}}^{(\text{III})}[\rho_1] \approx \mathcal{E}_{\text{co}}^{(\text{gas})}[\varrho_{1g}]_{(\varrho_{1g}=\rho_1)}, \quad (81)$$

where ϱ_{1g} is the spin-less, one-particle density-matrix of the Brueckner reference-state for an electron gas:

$$\tau_g(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{2} \varrho_{1g}(\mathbf{r}_1, \mathbf{r}_2) \delta_{\omega_1 \omega_2}. \quad (82)$$

For a *uniform* electron gas, Eq. (82) is denoted by

$$\tau_{\text{ug}}(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{2} \varrho_{1\text{ug}}(\mathbf{r}_1, \mathbf{r}_2) \delta_{\omega_1 \omega_2}, \quad (83)$$

and the density that is from the Brueckner, one-particle reference-state of a uniform electron gas, say $|\Theta_{\text{ug}}\rangle$, is given by

$$\varrho_{\text{ug}}(\mathbf{r}) = \varrho_{1\text{ug}}(\mathbf{r}, \mathbf{r}), \quad (84)$$

and this density is constant, i.e., it does not depend on $\mathbf{r}$; furthermore, we have

$$\varrho_{\text{ug}} = n_{\text{ug}}, \quad (85)$$

where n_{ug} is the density from the target state, say $|\Psi_{\text{ug}}\rangle$, of a uniform electron gas:

$$|\Psi_{\Theta_{\text{ug}}}^{(\eta)}\rangle = |\Psi_{\text{ug}}\rangle, \quad (86)$$

where we have used Eq. (24).

The correlation energy $\mathcal{E}_{\text{co}}^{(\text{ug})}$ of a uniform electron gas depends on the constant density n_{ug} ; it is given by the following relation:

$$\mathcal{E}_{\text{co}}^{(\text{ug})}(n_{\text{ug}}) = N \epsilon_{\text{co}}^{\text{ug}}(n_{\text{ug}}), \quad (87)$$

where $\epsilon_{\text{co}}^{\text{ug}}$ is the correlation-energy per particle and N is the number of particles. Using Eq. (85) gives

$$\mathcal{E}_{\text{co}}^{(\text{ug})}(\varrho_{\text{ug}}) = N \epsilon_{\text{co}}^{\text{ug}}(\varrho_{\text{ug}}). \quad (88)$$

Following Kohn and Sham [8], but using Eq. (88) instead of (87), we construct the following functional for an electron gas:

$$\mathcal{E}_{\text{co}}^{(\text{gas})}[\varrho_g] \approx \int \varrho_g(\mathbf{r}) \epsilon_{\text{co}}^{\text{ug}}(\varrho_g) d\mathbf{r}. \quad (89)$$

This functional can be expressed as one that appears to depend upon the one-particle density-matrix:

$$\mathcal{E}_{\text{co}}^{(\text{gas})}[\varrho_{1g}] \approx \iint \varrho_{1g}(\mathbf{r}_2, \mathbf{r}_1) \delta(\mathbf{r}_1 - \mathbf{r}_2) \epsilon_{\text{co}}^{\text{ug}}(\varrho_{1g}(\mathbf{r}_2, \mathbf{r}_1)) d\mathbf{r}_1 d\mathbf{r}_2, \quad (90)$$

where $(\varrho_{1g}(\mathbf{r}, \mathbf{r}) = \varrho_g(\mathbf{r}))$. Substituting the above expression into Eq. (81) gives

$$\bar{E}_{\text{co}}^{(\text{III})}[\rho_1] \approx \iint \rho_1(\mathbf{r}_2, \mathbf{r}_1) \delta(\mathbf{r}_1 - \mathbf{r}_2) \epsilon_{\text{co}}^{\text{ug}}(\rho_1(\mathbf{r}_2, \mathbf{r}_1)) d\mathbf{r}_1 d\mathbf{r}_2. \quad (91)$$

The correlation potential $\hat{\nu}_{\text{co}}^{\rho_1\text{I}^\text{II}}$ is defined by its kernel, given by Eq. (70). Because of the $\delta(\mathbf{r}_1 - \mathbf{r}_2)$ factor, $\hat{\nu}_{\text{co}}^{\rho_1\text{I}^\text{III}}$ is a local operator when using the above expressions for $\bar{E}_{\text{co}}^{(\text{III})}$; explicitly, using the results from Appendix A, we have the following approximate expressions:

$$\bar{E}_{\text{co}}^{(\text{III})}[\rho_1] \approx E_{\text{co}}^{\text{ug}}[\rho] = \int \rho(\mathbf{r}) \epsilon_{\text{co}}^{\text{ug}}(\rho) d\mathbf{r}, \quad (92)$$

$$\hat{\nu}_{\text{co}}^{\rho_1\text{I}^\text{III}}(\mathbf{r}) \approx \nu_{\text{co}}^{\text{ug}}(\mathbf{r}) = \frac{\delta E_{\text{co}}^{\text{ug}}[\rho]}{\delta \rho(\mathbf{r})}, \quad (93)$$

where, for future reference, these approximations are denoted by $E_{\text{co}}^{\text{ug}}$ and $\nu_{\text{co}}^{\text{ug}}$.

Using the exchange-energy formula by Dirac for a uniform electron gas, the exchange portion can be treated in an identical manner as the correlation; this approach yields the following local exchange-operator and exchange energy:

$$E_{\text{x}}[\rho_1] \approx E_{\text{x}}^{\text{ug}}[\rho] = \frac{3}{4} \left(\frac{3}{\pi} \right)^{1/3} \int \rho(\mathbf{r})^{4/3} d\mathbf{r}, \quad (94)$$

$$\hat{\nu}_{\text{x}}^{\rho_1}(\mathbf{r}) \approx \nu_{\text{x}}^{\text{ug}}(\mathbf{r}) = \frac{\delta E_{\text{x}}^{\text{ug}}[\rho]}{\delta \rho(\mathbf{r})} = \left(\frac{3}{\pi} \rho(\mathbf{r}) \right)^{1/3}, \quad (95)$$

where E_x^{ug} denotes the Dirac exchange-energy functional and v_x^{ug} is the corresponding exchange potential. The above four expressions are identical to the ones for the LDA of DFT, except that – in the current approach – ρ is *not* required to be the density of an exact wavefunction $|\Psi\rangle$; ρ is *only* required to be the density of a reference state $|\Phi\rangle$, and this is assured to be the case.

B. The Becke exchange correction

For closed-shell systems that are well described by a single determinant, it is well known that the Bruckner and Hartree–Fock reference-states, $|\Theta\rangle$ and $|\tilde{\Theta}\rangle$, are usually very similar. Hence, in these cases, the spinless, one-particle density-matrices from the Brueckner reference-state, ϱ_1 , is similar to the one from the Hartree–Fock reference state, $\tilde{\varrho}_1$. Therefore, an approximate exchange-functional $E_x[\rho_1]$ that depends on the density of the reference state, i.e.,

$$E_x[\rho_1] \approx \tilde{E}_x[\rho], \quad (96)$$

and is an accurate approximation with the Hartree–Fock approach, is also a reasonable approximation for the current approach. Of particular interest is the exchange functional obtained by Becke [57] – where this functional is an approximation of the Hartree–Fock exchange with empirical parameters fit to atomic systems; explicitly, this functional is given by

$$\tilde{E}_x[\rho] = E_x^{\text{ug}}[\rho] + \Delta E_x^{\text{B88}}[\rho], \quad (97)$$

where ΔE_x^{B88} is the Becke exchange correction to the uniform electron-gas term, E_x^{ug} . Since this correction term contains a gradient dependence, the general form of the functional can be written using the following integral relation:

$$\tilde{E}_x[\rho] = \int G_x(\rho, \nabla \rho) d\mathbf{r}. \quad (98)$$

By using Eqs. (96) and (97) and the following identity:

$$\nabla_1 \rho(\mathbf{r}_1) = [(\nabla_1 + \nabla_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1)]_{\mathbf{r}_2=\mathbf{r}_1}, \quad (99)$$

we obtain the following approximation:

$$E_x[\rho_1] \approx E_x^{\text{ug}}[\rho] + \Delta E_x^{\text{B88}}[\rho] = \iint \delta(\mathbf{r}_2 - \mathbf{r}_1) G_x \left(\rho_1(\mathbf{r}_2, \mathbf{r}_1), (\nabla_1 + \nabla_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1) \right) d\mathbf{r}_1 d\mathbf{r}_2.$$

Using this expression for the exchange energy, the exchange potential is local; using the result from Appendix A and the definition for v_x^{ug} given by Eq. (95), we have

$$\hat{v}_x^{\rho 1}(\mathbf{r}) \approx v_x^{\text{ug}}(\mathbf{r}) + v_x^{\text{B88}}(\mathbf{r}) = \frac{\delta E_x^{\text{ug}}[\rho]}{\delta \rho(\mathbf{r})} + \frac{\delta \Delta E_x^{\text{B88}}[\rho]}{\delta \rho(\mathbf{r})}, \quad (100)$$

where this expression defines $v_x^{\text{B88}}(\mathbf{r})$.

C. The LYP Functional

Consider the approximation given by Eq. (78), where ($\eta = \text{III}$), and the correlation energy $\mathcal{E}_{\text{co}}$ – which is also a functional – is taken to be the one from the helium atom:

$$\bar{E}_{\text{co}}^{(\text{III})}[\gamma, v] = \mathcal{E}_{\text{co}}[\tau_{\text{He}}, v_{\text{He}}]_{(\tau_{\text{He}}=\gamma, v_{\text{He}}=v)}, \quad (101)$$

where τ_{He} is the Brueckner, one-particle, density-matrix for the helium atom, and v_{He} is the external potential for this system. Furthermore, since the Brueckner density-matrix τ_{He} is approximately equal to the Hartree–Fock one, say $\tilde{\tau}_{\text{He}}$, we can write

$$E_{\text{co}}^{(\text{III})}[\gamma, v] \approx \mathcal{E}_{\text{co}}[\tilde{\tau}_{\text{He}}, v_{\text{He}}]_{(\tilde{\tau}_{\text{He}}=\gamma, v_{\text{He}}=v)}. \quad (102)$$

The dominant portion of the correlation energy comes from the electron-repulsion contribution. The portion arising from the external potential is smaller; since v is a one-body operator, this term is probably treated well as part of the first-order energy E_1 . Therefore, neglecting to make the substitution for the potential, i.e., ($v_{\text{He}} = v$), should only yield a small error:

$$E_{\text{co}}^{(\text{III})}[\gamma] \approx \mathcal{E}_{\text{co}}[\tilde{\tau}_{\text{He}}, v_{\text{He}}]_{(\tilde{\tau}_{\text{He}}=\gamma)}. \quad (103)$$

A well known approximation for $\mathcal{E}_{\text{co}}[\tilde{\tau}_{\text{He}}, v_{\text{He}}]$ – that is valid for other systems also – is given by Colle and Salvetti functional [14, 15]:

$$\mathcal{E}_{\text{co}}[\tilde{\tau}_{\text{He}}, v_{\text{He}}] \approx \mathcal{E}_{\text{co}}^{\text{CS}}[\tilde{\tau}_{\text{He}}]. \quad (104)$$

The above two relations suggest the following approximation:

$$E_{\text{co}}^{(\text{III})}[\gamma] \approx \mathcal{E}_{\text{co}}^{\text{CS}}[\tilde{\tau}_{\text{He}}]_{(\tilde{\tau}_{\text{He}}=\gamma)}. \quad (105)$$

While this is probably a reasonable approximation, we do have some reservations about it. In particular, a universal functional is used to approximate one that depends on the external

potential. This violates, at least, the spirit of the method. We would therefore prefer to use the approximation given by Eq. (101) or (102). Unfortunately, however, no such functionals are available. Therefore, let us proceed.

For a closed-shell system, Eq. (105) becomes

$$\bar{E}_{\text{co}}^{(\text{III})}[\rho_1] \approx E_{\text{co}}^{\text{CS}}[\rho_1] = \mathcal{E}_{\text{co}}^{\text{CS}}[\tilde{\varrho}_{\text{He}}]_{(\tilde{\varrho}_{\text{He}}=\rho_1)}, \quad (106)$$

where $\tilde{\varrho}_{\text{He}}$ is the Hartree–Fock spin-less one-particle density-matrix for the helium atom, and, for future reference, this approximation is denoted by $E_{\text{co}}^{\text{CS}}[\rho_1]$. An expression for $E_{\text{co}}^{\text{CS}}[\rho_1]$ is derived by Lee, Parr, and Yang [14] that is equivalent to the one derived by Colle–Salvetti; it has the following form:

$$E_{\text{co}}^{\text{CS}}[\rho_1] = \int G_{\text{co}} \left(\rho_1(\mathbf{r}_1, \mathbf{r}_1), \nabla_1 \rho_1(\mathbf{r}_1, \mathbf{r}_1), \nabla_1^2 \rho_1(\mathbf{r}_1, \mathbf{r}_1), (\nabla_1 \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1))_{\mathbf{r}_2=\mathbf{r}_1} \right) d\mathbf{r}_1, \quad (107)$$

and using this Eq. and the previous one, we have

$$\bar{E}_{\text{co}}^{(\text{III})}[\rho_1] \approx \iint \delta(\mathbf{r}_2 - \mathbf{r}_1) G_{\text{co}} \left(\rho_1(\mathbf{r}_1, \mathbf{r}_1), \nabla_1 \rho_1(\mathbf{r}_1, \mathbf{r}_1), \nabla_1^2 \rho_1(\mathbf{r}_1, \mathbf{r}_1), \nabla_1 \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1) \right) d\mathbf{r}_1 d\mathbf{r}_2. \quad (108)$$

When the functional derivative is taken to determine the kernel of the correlation potential $\nu_{\text{co}}^{\rho_1 \text{III}}(\mathbf{r}_1, \mathbf{r}_2)$ – as defined by Eq. (70) – and $\bar{E}_{\text{co}}^{(\text{III})}$ is approximated by the above relation, the $\delta(\mathbf{r}_2 - \mathbf{r}_1)$ term survives; the correlation potential $\hat{\nu}_{\text{co}}^{\rho_1 \text{III}}$ is a local operator. Denoting this approximation by $\nu_{\text{co}}^{\text{CS}}(\mathbf{r})$, we have

$$\hat{\nu}_{\text{co}}^{\rho_1 \text{III}}(\mathbf{r}) \approx \nu_{\text{co}}^{\text{CS}}(\mathbf{r}). \quad (109)$$

Using the exchange potential given by Eq. (100) – or any other local exchange potential – and the above local correlation operator, the exchange-correlation potential $\hat{\nu}_{\text{xc}}^{\rho_1 \eta}$ – given by Eq. (71) – is also local. Therefore, the reference-state $|\Phi\rangle$ obtained from the following non-interacting Hamiltonian:

$$H_s^{\rho_1} = \sum_i \left(-\frac{1}{2} \nabla_i^2 \right) + \sum_i \nu_s^{\rho_1}(\mathbf{r}_i), \quad (110)$$

is also obtained from a local potential, where

$$\nu_s^{\rho_1}(\mathbf{r}_1) = v(\mathbf{r}_1) + \int r_{12}^{-1} \varrho_1(\mathbf{r}_2) d\mathbf{r}_2 + v_{\text{x}}^{\text{ug}}(\mathbf{r}) + v_{\text{x}}^{\text{B88}}(\mathbf{r}_1) + v_{\text{co}}^{\text{CS}}(\mathbf{r}_1), \quad (111)$$

and Eq. (66), for the occupied orbitals, becomes

$$\left(-\frac{1}{2}\nabla_1^2 + \nu_s^{e_1}(\mathbf{r}_1)\right) \chi_w^{e_1}(\mathbf{r}_1) = \xi_w^{e_1} \chi_w^{e_1}(\mathbf{r}_1). \quad (112)$$

If these equations are solved in a self consistent (SCF) manner, a series of spin-less, one-particle density-matrices are generated: $\rho_1^0, \rho_1^1, \rho_1^2, \dots$; for the n th iteration, we have

$$\left(-\frac{1}{2}\nabla_1^2 + \nu_s^{\rho_1^n}(\mathbf{r}_1)\right) \chi_w^{\rho_1^n}(\mathbf{r}_1) = \xi_w^{\rho_1^n} \chi_w^{\rho_1^n}(\mathbf{r}_1), \quad (113)$$

and this iteration yields a non-interacting state, say $|\Phi^{n+1}\rangle$, with a one-particle density-matrix of ρ_1^{n+1} ; in addition, $|\Phi^{n+1}\rangle$ is an eigenfunction of a non-interaction Hamiltonian, $H_s^{\rho_1^n}$. From the Hohenberg and Kohn theorem [58], each non-interacting states generated in the SCF procedure are unique functions of their electron densities. e.g., $|\Phi^n(\rho)\rangle$; furthermore, because of the one-to-one correspondence between one-particle density-matrices and non-interacting states, all one-particle density-matrices generated from the SCF procedure are determined by their density, i.e., $\rho_1(\rho)$. And since for an SCF computation, these are the only one-particle density-matrices that are required, we can write

$$E_{\text{co}}^{\text{cs}}[\rho_1] = E_{\text{co}}^{\text{cs}}[\rho]. \quad (114)$$

Because of the general form of $E_{\text{co}}^{\text{cs}}[\rho_1]$ given by Eq. (107) we need only an expression for $(\nabla_1 \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1))_{\mathbf{r}_2=\mathbf{r}_1}$. Following Lee, Parr, and Yang, we use the following approximation:

$$\left(\nabla_1 \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1)\right)_{\mathbf{r}_2=\mathbf{r}_1} \approx \frac{3}{5}(3\pi^2)^{2/3} \rho(\mathbf{r}_1)^{5/3} + \frac{1}{36} \frac{|\nabla_1 \rho(\mathbf{r}_1)|^2}{\rho(\mathbf{r}_1)} + \frac{1}{4} \nabla_1^2 \rho(\mathbf{r}_1), \quad (115)$$

and we have

$$E_{\text{co}}^{\text{cs}}[\rho] \approx E_{\text{co}}^{\text{lyp}}[\rho], \quad (116)$$

where $E_{\text{co}}^{\text{lyp}}[\rho]$ is the LYP functional obtained by substituting the approximation given by Eq. (115) into the $E_{\text{co}}^{\text{cs}}[\rho_1]$ functional. Using the general of form of $E_{\text{co}}^{\text{cs}}[\rho_1]$ given by Eq. (107), we have

$$E_{\text{co}}^{\text{lyp}}[\rho] = \int G_{\text{co}}\left(\rho(\mathbf{r}), \nabla \rho(\mathbf{r}), \nabla^2 \rho(\mathbf{r})\right) d\mathbf{r}. \quad (117)$$

Using integration by parts, Miehlich, Savin, Stoll, and Preuss, eliminate the Laplacian terms and obtain the general form given by [59]

$$E_{\text{co}}^{\text{lyp}}[\rho] = \int G_{\text{co}}\left(\rho(\mathbf{r}), \nabla \rho(\mathbf{r})\right) d\mathbf{r}. \quad (118)$$

Using Eqs. (106), (114), (116), (118), and (99) we obtain the following approximation for the correlation-energy functional:

$$\bar{E}_{\text{co}}^{(\text{III})}[\rho_1] \approx E_{\text{co}}^{\text{lyp}}[\rho] = \iint \delta(\mathbf{r}_2 - \mathbf{r}_1) G_{\text{co}} \left(\rho_1(\mathbf{r}_2, \mathbf{r}_1), (\nabla_1 + \nabla_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1) \right) d\mathbf{r}_1 d\mathbf{r}_2, \quad (119)$$

and using the results from Appendix A, the local correlation potential is given by

$$\hat{\nu}_{\text{co}}^{\rho_1^{\text{III}}}(\mathbf{r}) \approx \nu_{\text{co}}^{\text{LYP}}(\mathbf{r}) = \frac{\delta E_{\text{co}}^{\text{lyp}}[\rho]}{\delta \rho(\mathbf{r})}. \quad (120)$$

Using these two approximations for $E_{\text{co}}^{\text{lyp}}[\rho]$ and $\hat{\nu}_{\text{co}}^{\rho_1^{\text{III}}}(\mathbf{r})$ yields an identical approach as the LYP method. If, in addition, the Becke exchange functional, given by Eq. (100), is used, the method is equivalent to the one known as BLYP.

D. The B3LYP Functional

Using Eq. (43b), and the uniform-electron-gas approximation for the correlation-energy functional, Eq. (92), we obtain an approximate exchange-correlation functional, given by

$$\bar{E}_{\text{xc}}^{(\text{III})}[\rho_1] \approx E_{\text{x}}[\rho_1] + E_{\text{co}}^{\text{ug}}[\rho]. \quad (121)$$

An alternative approximation is given by the BLYP functional that uses Eqs. (100) and (119):

$$\bar{E}_{\text{xc}}^{(\text{III})}[\rho_1] \approx E_{\text{x}}^{\text{ug}}[\rho] + \Delta E_{\text{x}}^{\text{B88}}[\rho] + E_{\text{co}}^{\text{lyp}}[\rho]. \quad (122)$$

If both of these approximations are reasonable, then so is a linear combination of the two:

$$\bar{E}_{\text{xc}}^{(\text{III})}[\rho_1] \approx b \left(E_{\text{x}}^{\text{ug}}[\rho] + \Delta E_{\text{x}}^{\text{B88}}[\rho] + E_{\text{co}}^{\text{lyp}}[\rho] \right) + (1 - b) \left(E_{\text{x}}[\rho_1] + E_{\text{co}}^{\text{ug}}[\rho] \right), \quad (123)$$

where the semi-empirical parameter b is at our disposal. Using the approximation for $E_{\text{co}}^{\text{ug}}$, given by Vosko, Wilk, and Nusair [60], denoted by $E_{\text{co}}^{\text{vwn}}$, we have

$$\bar{E}_{\text{xc}}^{(\text{III})}[\rho_1] \approx b \left(E_{\text{x}}^{\text{ug}}[\rho] + \Delta E_{\text{x}}^{\text{B88}}[\rho] + E_{\text{co}}^{\text{lyp}}[\rho] \right) + (1 - b) \left(E_{\text{x}}[\rho_1] + E_{\text{co}}^{\text{vwn}}[\rho] \right). \quad (124)$$

Insertion of additional empirical parameters that deviate slightly from unity, we have

$$\bar{E}_{\text{xc}}^{(\text{III})}[\rho_1] \approx b \left(E_{\text{x}}^{\text{ug}}[\rho] + \alpha \Delta E_{\text{x}}^{\text{B88}}[\rho] + \beta E_{\text{co}}^{\text{lyp}}[\rho] \right) + (1 - b) \left(E_{\text{x}}[\rho_1] + \gamma E_{\text{co}}^{\text{vwn}}[\rho] \right), \quad (125)$$

where we can reduce the four variables to three by introducing a constraint, e.g., $(\gamma[b, \beta])$. By choosing $(b = 0.80)$, $(\alpha = 0.90)$, $(\beta = 1.01)$, $(\gamma = (1 - \beta b)(1 - b)^{-1} = 0.95)$ we get the same exchange-correlation energy functional as the one employed in the B3LYP approach. In particular, we have $(b = 1 - a_0)$, $(b\alpha = a_x)$, and $(b\beta = a_c)$, where a_0 , a_x , and a_c , are 0.20, 0.72, and 0.81, respectively.

V. ACKNOWLEDGMENTS

The author thanks Kimihiko Hirao for useful discussions.

APPENDIX A: DERIVATION OF SOME IDENTITIES

Consider the following density functional:

$$E_i[\rho] = \int G_i(\rho) d\mathbf{r}, \quad (\text{A1})$$

and a corresponding potential, given by [2, 7, 61]

$$v_i(\mathbf{r}) = \frac{\delta E_i[\rho]}{\delta \rho(\mathbf{r})} = \frac{\partial G_i(\rho)}{\partial \rho(\mathbf{r})}. \quad (\text{A2})$$

If the integrand $G_i(\rho)$ does not contain gradient terms, we can express the density functional as one that appears to depend on the spin-less one-particle density-matrix:

$$E_i[\rho] = E_i[\rho_1] = \iint \delta(\mathbf{r}_1 - \mathbf{r}_2) G_i(\rho_1) d\mathbf{r}_1 d\mathbf{r}_2, \quad (\text{A3})$$

and its corresponding operator, denoted by $\hat{v}_i$, is defined by its kernel, given by

$$v_i(\mathbf{r}_1, \mathbf{r}_2) = \frac{\delta E_i[\rho_1]}{\delta \rho_1(\mathbf{r}_2, \mathbf{r}_1)} = \delta(\mathbf{r}_1 - \mathbf{r}_2) \frac{\partial G_i(\rho_1)}{\partial \rho_1(\mathbf{r}_2, \mathbf{r}_1)}. \quad (\text{A4})$$

Operating with $\hat{v}_i$ upon an arbitrary function, ϕ , and using Eqs. (A2) and (A4), we obtain the following:

$$\begin{aligned} \hat{v}_i \phi(\mathbf{r}_1) &= \int v_i(\mathbf{r}_1, \mathbf{r}_2) \phi(\mathbf{r}_2) d\mathbf{r}_2 = \int \delta(\mathbf{r}_1 - \mathbf{r}_2) \frac{\partial G_i(\rho_1(\mathbf{r}_2, \mathbf{r}_1))}{\partial \rho_1(\mathbf{r}_2, \mathbf{r}_1)} \phi(\mathbf{r}_2) d\mathbf{r}_2 \\ &= \frac{\partial G_i(\rho_1(\mathbf{r}_1, \mathbf{r}_1))}{\partial \rho_1(\mathbf{r}_1, \mathbf{r}_1)} \phi(\mathbf{r}_1) = \frac{\partial G_i(\rho(\mathbf{r}_1))}{\partial \rho(\mathbf{r}_1)} \phi(\mathbf{r}_1) = v_i(\mathbf{r}_1) \phi(\mathbf{r}_1). \end{aligned} \quad (\text{A5})$$

Hence, the potential and operator are identical:

$$\hat{v}_i = v_i(\mathbf{r}). \quad (\text{A6})$$

Now let the energy functional have a gradient dependence:

$$E_i[\rho] = \int G_i(\rho, \nabla \rho) d\mathbf{r}, \quad (\text{A7})$$

where its corresponding potential is given by [2, 7, 61]

$$v_i(\mathbf{r}) = \frac{\delta E_i[\rho]}{\delta \rho(\mathbf{r})} = \frac{\partial G_i(\rho)}{\partial \rho(\mathbf{r})} - \frac{d}{d\mathbf{r}} \cdot \frac{\partial G_i(\rho)}{\partial \nabla \rho(\mathbf{r})}. \quad (\text{A8})$$

Using the identity (99), we can express the density functional as a one-particle density-matrix functional:

$$E_i[\rho] = E_i[\rho_1] = \iint \delta(\mathbf{r}_1 - \mathbf{r}_2) G_i \left(\rho_1(\mathbf{r}_2, \mathbf{r}_1), (\nabla_1 + \nabla_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1) \right) d\mathbf{r}_1 d\mathbf{r}_2, \quad (\text{A9})$$

and its corresponding operator, denoted by $\hat{v}_i$, is defined by its kernel, given by

$$\begin{aligned} v_i(\mathbf{r}_1, \mathbf{r}_2) &= \frac{\delta E_i[\rho_1]}{\delta \rho(\mathbf{r}_2, \mathbf{r}_1)} \\ &= \delta(\mathbf{r}_1 - \mathbf{r}_2) \left(\frac{\partial G_i(\rho_1)}{\partial \rho_1(\mathbf{r}_2, \mathbf{r}_1)} - \frac{d}{d\mathbf{r}_1} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_1 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} - \frac{d}{d\mathbf{r}_2} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} \right). \end{aligned} \quad (\text{A10})$$

Operating with $\hat{v}_i$ upon an arbitrary function ϕ , using Eqs. (A10) and (A8), and an identity, given by

$$\left(\frac{d}{d\mathbf{r}_1} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_1 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} + \frac{d}{d\mathbf{r}_2} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} \right)_{\mathbf{r}_2=\mathbf{r}_1} = \frac{d}{d\mathbf{r}_1} \cdot \frac{\partial G_i(\rho)}{\partial \nabla_1 \rho(\mathbf{r}_1)}, \quad (\text{A11})$$

we obtain the following:

$$\begin{aligned} \hat{v}_i \phi(\mathbf{r}_1) &= \int v_i(\mathbf{r}_1, \mathbf{r}_2) \phi(\mathbf{r}_2) d\mathbf{r}_2 \\ &= \int \delta(\mathbf{r}_1 - \mathbf{r}_2) \left(\frac{\partial G_i(\rho_1)}{\partial \rho_1(\mathbf{r}_2, \mathbf{r}_1)} - \frac{d}{d\mathbf{r}_1} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_1 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} - \frac{d}{d\mathbf{r}_2} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} \right) \phi(\mathbf{r}_2) d\mathbf{r}_2 \\ &= \left[\frac{\partial G_i(\rho(\mathbf{r}_1))}{\partial \rho(\mathbf{r}_1)} - \left(\frac{d}{d\mathbf{r}_1} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_1 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} + \frac{d}{d\mathbf{r}_2} \cdot \frac{\partial G_i(\rho_1)}{\partial \nabla_2 \rho_1(\mathbf{r}_2, \mathbf{r}_1)} \right)_{\mathbf{r}_2=\mathbf{r}_1} \right] \phi(\mathbf{r}_1) \\ &= \left(\frac{\partial G_i(\rho(\mathbf{r}_1))}{\partial \rho(\mathbf{r}_1)} - \frac{d}{d\mathbf{r}_1} \cdot \frac{\partial G_i(\rho)}{\partial \nabla_1 \rho(\mathbf{r}_1)} \right) \phi(\mathbf{r}_1) = v_i(\mathbf{r}_1) \phi(\mathbf{r}_1). \end{aligned} \quad (\text{A12})$$

Hence, we have

$$\hat{v}_i = v_i(\mathbf{r}). \quad (\text{A13})$$

-
- [1] R. M. Dreizler and E. K. U. Gross, *Density Functional Theory: An Approach to the Quantum Many-Body Problem* (Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, Hong Kong, Barcelona, 1990).
- [2] R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules* (Oxford University Press, New York, 1989).
- [3] M. Springborg, ed., *Density Functional Methods in Chemistry and Material Science* (Wiley, Chichester, 1997).
- [4] D. E. Ellis, ed., *Density Functional Methods of Molecules, Clusters, and Solids* (Kluwer, Dordrecht, 1995).
- [5] E. K. U. Gross and R. M. Dreizler, eds., *Density Functional Theory* (Plenum, New York, 1994).
- [6] J. Seminario and P. Politzer, eds., *Modern Density Functional Theory. A Tool for Chemistry* (Elsevier, Amsterdam, 1995).
- [7] N. C. Handy, in *Europe Summerschool in Quantum Chemistry*, B. O. Roos and P. -O. Widmark eds. (1997), 2nd ed., (see internet address www.teokem.lu.se/esqc).
- [8] W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
- [9] A. D. Becke, J. Chem. Phys. **98**, 5648 (1993).
- [10] K. Burke, M. Ernzerhof, and J. P. Perdew, Chem. Phys. Lett. **265**, 115 (1997).
- [11] J. P. Perdew, M. Ernzerhof, and K. Burke, J. Chem. Phys. **105**, 9982 (1996).
- [12] M. Ernzerhof, Chem. Phys. Lett. **263**, 499 (1996).
- [13] P. J. Stephens, J. F. Devlin, and C. F. Chabalowski, J. Phys. Chem. **98**, 11623 (1994), (see also internet address <http://www.gaussian.com/q3.htm>).
- [14] C. Lee, W. Yang, and R. G. Parr, Phys. Rev. B **37**, 785 (1988).
- [15] R. Colle and O. Salvetti, Theor. Chim. Acta **37**, 329 (1975).
- [16] J. P. Finley, arXiv:physics/0308056 (August 14, 2003).
- [17] J. P. Finley, Phys. Rev. A. (2003), (submitted).
- [18] I. Lindgren and J. Morrison, *Atomic Many-Body Theory*, Springer Series on Atoms and Plasmas (Springer-Verlag, New York, Berlin, Heidelberg, 1986), 2nd ed.
- [19] F. E. Harris, H. J. Monkhorst, and D. L. Freeman, *Algebraic and Diagrammatic Methods in*

- Many-Fermion Theory* (Oxford University Press, New York, 1992).
- [20] I. Lindgren and S. Solomonson, *Int. J. Quantum Chem.* **90**, 294 (2002).
 - [21] A. Seidl, A. Görling, P. Vogl, J. A. Majewski, and M. Levy, *Phys. Rev. B* **53**, 3764 (1996).
 - [22] A. Szabo and N. S. Ostlund, *Modern Quantum Chemistry: Introduction to Advanced Electronic Structure Theory* (Macmillan, New York, 1982).
 - [23] J.-P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems* (MIT Press, Cambridge, Mass., 1986).
 - [24] P. A. M. Dirac, *Proc. Cambridge Phil. Soc.* **26**, 376 (1930).
 - [25] P. A. M. Dirac, *Proc. Cambridge Phil. Soc.* **27**, 240 (1931).
 - [26] P. -O. Löwdin, *Phys.* **97**, 1474 (1955).
 - [27] P. -O. Löwdin, *Phys.* **97**, 1490 (1955).
 - [28] R. McWeeny, *Rev. Mod. Phys.* **32**, 335 (1960).
 - [29] K. A. Brueckner, *Phys. Rev.* **96**, 508 (96), (See Ref. 4 for more citations from Brueckner and coworkers.).
 - [30] R. K. Nesbet, *Phys. Rev.* **109**, 1632 (1958).
 - [31] W. Brenig, *Nucl. Phys.* **22**, 177 (1961).
 - [32] P. O. Löwdin, *J. Math. Phys.* **3**, 1171 (1962).
 - [33] W. Kutzelnigg and V. H. Smith, *J. Chem. Phys.* **41**, 896 (1964).
 - [34] J. Paldus and J. Čížek, *Phys. Scripta* **21**, 251 (1980).
 - [35] R. A. Chiles and C. E. Dykstra, *J. Chem. Phys.* **74**, 4544 (1981).
 - [36] L. Z. Stolarczyk and H. J. Monkhorst, *Int. J. Quantum Chem.* **18**, 267 (1984).
 - [37] N. C. Handy, J. A. Pople, M. Head-Gordon, K. Raghavachari, and G. W. Trucks, *Chem. Phys. Lett.* **164**, 185 (1985).
 - [38] N. C. Handy, J. A. Pople, M. Head-Gordon, K. Raghavachari, and G. W. Trucks, *Chem. Phys. Lett.* **164**, 185 (1989).
 - [39] K. K. Raghavachari, J. A. Pople, E. S. Replogle, M. Head-Gordon, and N. C. Handy, *Chem. Phys. Lett.* **167**, 115 (1990).
 - [40] K. Hirao, in *Self Consistent Field Theory, Studies in Physical and Theoretical Chemistry* (Elsevier, Amsterdam, 1990), vol. 90.
 - [41] J. F. Stanton, J. Gauss, and R. J. Bartlett, *J. Chem. Phys.* **97**, 5554 (1992).
 - [42] C. Hampel, K. A. Peterson, and H. -J. Werner, *Chem. Phys. Lett.* **190**, 1 (1992).

- [43] G. E. Scuseria, Chem. Phys. Lett. **226**, 251 (1994).
- [44] D. H. Kobe, Nucl. Phys. **3**, 417 (1971).
- [45] L. Schäfer and H. A. Weidenmüller, Nucl. Phys. **A174**, 1 (1971).
- [46] J. Hubbard, Proc. Roy. Soc. London **A 240**, 539 (1957).
- [47] F. Coester, Nucl. Phys. **7**, 421 (1958).
- [48] J. Čížek, J. Chem. Phys. **45**, 4256 (1966).
- [49] J. Čížek, Adv. Chem. Phys. **14**, 35 (1969).
- [50] J. Čížek and J. Paldus, Int. J. Quantum Chem. **5**, 359 (1971).
- [51] I. Lindgren, Int. J. Quantum Chem. **S12**, 33 (1978).
- [52] R. J. Bartlett and G. D. Purvis, Int. J. Quantum Chem. **14**, 561 (1978).
- [53] J. A. Pople, R. Krishnan, H. B. Schlegel, and J. S. Binkley, Int. J. Quantum Chem. **S14**, 545 (1978).
- [54] H.f Schaefer III, *Electronic Structures of Atoms and Molecules—A Survey of Rigorous Quantum Mechanical Methods* (Addison-Wesley, Reading, 1972).
- [55] J. P. Finley (2003), (to be published).
- [56] I. Lindgren, J. Phys. B **7**, 2441 (1974).
- [57] A. D. Becke, Phys. Rev. A **38**, 3098 (1988).
- [58] P. Hohenberg and W. Kohn, Phys. Rev. **136B**, 864 (1964).
- [59] B. Miehloch, A. S. abd H. Stoll, and H. Preuss, Chem. Phys. Lett. **157**, 200 (1989).
- [60] S. H. Vosko, L. Wilk, and M. Nusair, Can. J. Phys. **58**, 1200 (1980).
- [61] I. M. Gelfand and S. Fomin, *Calculus of variations* (1963).